

# Bipartite graphs with uniquely restricted maximum matchings and their corresponding greedoids

Vadim E. Levit and Eugen Mandrescu  
 Department of Computer Science  
 Holon Academic Institute of Technology  
 52 Golomb Str., P.O. Box 305  
 Holon 58102, ISRAEL  
 {levitv, eugen\_m}@barley.cteh.ac.il

## Abstract

A *maximum stable set* in a graph  $G$  is a stable set of maximum size.  $S$  is a *local maximum stable set* of  $G$ , and we write  $S \in \Psi(G)$ , if  $S$  is a maximum stable set of the subgraph spanned by  $S \cup N(S)$ , where  $N(S)$  is the neighborhood of  $S$ . A matching  $M$  is *uniquely restricted* if its saturated vertices induce a subgraph which has a unique perfect matching, namely  $M$  itself. Nemhauser and Trotter Jr. [12], proved that any  $S \in \Psi(G)$  is a subset of a maximum stable set of  $G$ . In [10] we have shown that the family  $\Psi(T)$  of a forest  $T$  forms a greedoid on its vertex set. In this paper we demonstrate that for a bipartite graph  $G$ ,  $\Psi(G)$  is a greedoid on its vertex set if and only if all its maximum matchings are uniquely restricted.

## 1 Introduction

Throughout this paper  $G = (V, E)$  is a simple (i.e., a finite, undirected, loopless and without multiple edges) graph with vertex set  $V = V(G)$  and edge set  $E = E(G)$ . If  $X \subset V$ , then  $G[X]$  is the subgraph of  $G$  spanned by  $X$ . By  $G - W$  we mean the subgraph  $G[V - W]$ , if  $W \subset V(G)$ . We also denote by  $G - F$  the partial subgraph of  $G$  obtained by deleting the edges of  $F$ , for  $F \subset E(G)$ , and we write shortly  $G - e$ , whenever  $F = \{e\}$ . If  $X, Y \subset V$  are disjoint and non-empty, then by  $(X, Y)$  we mean the set  $\{xy : xy \in E, x \in X, y \in Y\}$ . The *neighborhood* of a vertex  $v \in V$  is the set  $N(v) = \{w : w \in V \text{ and } vw \in E\}$ . If  $|N(v)| = 1$ , then  $v$  is a *pendant vertex* of  $G$ ; by  $\text{pend}(G)$  we designate the set of all pendant vertices of  $G$ . We denote the *neighborhood* of  $A \subset V$  by  $N_G(A) = \{v \in V - A : N(v) \cap A \neq \emptyset\}$  and its *closed neighborhood* by  $N_G[A] = A \cup N(A)$ , or shortly,  $N(A)$  and  $N[A]$ , if no ambiguity.  $K_n, C_n$  denote respectively, the complete graph on  $n \geq 1$  vertices and the chordless cycle on  $n \geq 3$  vertices. By  $G = (A, B, E)$  we mean a bipartite graph having  $\{A, B\}$  as its standard bipartition.

A *stable* set in  $G$  is a set of pairwise non-adjacent vertices. A stable set of maximum size will be referred to as a *maximum stable set* of  $G$ , and the *stability number* of  $G$ , denoted by  $\alpha(G)$ , is the cardinality of a maximum stable set in  $G$ . Let  $\Omega(G)$  stand for the set of all maximum stable sets of  $G$ . A set  $A \subseteq V(G)$  is a *local maximum stable set* of  $G$  if  $A$  is a maximum stable set in the subgraph spanned by  $N[A]$ , i.e.,  $A \in \Omega(G[N[A]])$ , [10]. In the sequel, by  $\Psi(G)$  we denote the set of all local maximum stable sets of the graph  $G$ . For instance, any set  $S \subseteq \text{pend}(G)$  belongs to  $\Psi(G)$ , while the converse is not generally true; e.g.,  $\{a\}, \{e, d\} \in \Psi(G)$  and  $\{e, d\} \cap \text{pend}(G) = \emptyset$ , where  $G$  is the graph in Figure 1.




Figure 1: A graph with diverse local maximum stable sets.

Not any stable set of a graph  $G$  is included in some maximum stable set of  $G$ . For example, there is no  $S \in \Omega(G)$  such that  $\{c, f\} \subset S$ , where  $G$  is the graph depicted in Figure 4. The following theorem due to Nemhauser and Trotter Jr. [12], shows that some special maximum stable sets can be enlarged to maximum stable sets.

**Theorem 1.1** [12] *Any local maximum stable set of a graph is a subset of a maximum stable set.*

Let us notice that the converse of Theorem 1.1 is not generally true. For instance,  $C_n$ ,  $n \geq 4$ , has no proper local maximum stable set. The graph  $G$  in Figure 1 shows another counterexample: any  $S \in \Omega(G)$  contains some local maximum stable set, but these local maximum stable sets are of different cardinalities. As examples,  $\{a, d, f\} \in \Omega(G)$  and  $\{a\}, \{d, f\} \in \Psi(G)$ , while for  $\{b, e, g\} \in \Omega(G)$  only  $\{e, g\} \in \Psi(G)$ .

In [10] we have proved the following result:

**Theorem 1.2** *The family of local maximum stable sets of a forest of order at least two forms a greedoid on its vertex set.*

Theorem 1.2 is not specific for forests. For instance, the family  $\Psi(G)$  of the graph  $G$  in Figure 2 is a greedoid.




Figure 2: A graph whose family of local maximum stable sets forms a greedoid.

The definition of greedoids we use in the sequel is as follows.

**Definition 1.3** [1], [6] A *greedoid* is a pair  $(E, \mathcal{F})$ , where  $\mathcal{F} \subseteq 2^E$  is a set system satisfying the following conditions:

(Accessibility) for every non-empty  $X \in \mathcal{F}$  there is an  $x \in X$  such that  $X - \{x\} \in \mathcal{F}$ ;  
(Exchange) for  $X, Y \in \mathcal{F}$ ,  $|X| = |Y| + 1$ , there is an  $x \in X - Y$  such that  $Y \cup \{x\} \in \mathcal{F}$ .

Clearly,  $\Omega(G) \subseteq \Psi(G)$  holds for any graph  $G$ . It is worth observing that if  $\Psi(G)$  is a greedoid and  $S \in \Psi(G)$ ,  $|S| = k \geq 2$ , then by accessibility property, there is a chain

$$\{x_1\} \subset \{x_1, x_2\} \subset \dots \subset \{x_1, \dots, x_{k-1}\} \subset \{x_1, \dots, x_{k-1}, x_k\} = S$$

such that  $\{x_1, x_2, \dots, x_j\} \in \Psi(G)$ , for all  $j \in \{1, \dots, k-1\}$ . Such a chain we call an *accessibility chain* for  $S$ . As an example, for  $S = \{a, c, e\} \in \Psi(G)$ , where  $G$  is the graph in Figure 2, an accessibility chain is  $\{a\} \subset \{a, e\} \subset S$ .

A *matching* in a graph  $G = (V, E)$  is a set of edges  $M \subseteq E$  having the property that no two edges of  $M$  share a common vertex. We denote the size of a *maximum matching* (a matching of maximum cardinality) by  $\mu(G)$ . A *perfect matching* is a matching saturating all the vertices of the graph.

Let us recall that  $G$  is a *König-Egerváry graph* provided  $\alpha(G) + \mu(G) = |V(G)|$ , [2], [7]. As a well-known example, any bipartite graph is a König-Egerváry graph. Some non-bipartite König-Egerváry graphs are presented in Figure 7.

A matching  $M = \{a_i b_i : a_i, b_i \in V(G), 1 \leq i \leq k\}$  of a graph  $G$  is called a *uniquely restricted matching* if  $M$  is the unique perfect matching of the subgraph  $G[\{a_i, b_i : 1 \leq i \leq k\}]$ , [4] (first time this kind of matching appeared in [5] for bipartite graphs under the name "*constrained matching*"). Let  $\mu_r(G)$  be the maximum size of a uniquely restricted matching in  $G$ . Clearly,  $0 \leq \mu_r(G) \leq \mu(G)$  holds for any graph  $G$ . For instance,  $0 = \mu_r(C_{2n}) < n = \mu(C_{2n})$ , while  $\mu_r(C_{2n+1}) = \mu(C_{2n+1}) = n$ .

In this paper we characterize the bipartite graphs whose family of local maximum stable sets are greedoids. Namely, we prove that for a bipartite graph  $G$ , the family  $\Psi(G)$  is a greedoid on the vertex set of  $G$  if and only if all its maximum matchings are uniquely restricted.

Golumbic, Hirst and Lewenstein have shown in [4] that  $\mu_r(G) = \mu(G)$  holds when  $G$  is a tree or has only odd cycles. Our findings reveal another class of graphs enjoying this property.

## 2 Preliminary results

An edge  $e$  of a graph  $G$  is  $\alpha$ -critical ( $\mu$ -critical) if  $\alpha(G) < \alpha(G - e)$  ( $\mu(G) > \mu(G - e)$ , respectively). Let us observe that there is no general connection between the  $\alpha$ -critical and the  $\mu$ -critical edges of a graph. For instance, the edge  $e$  of the graph  $G_1$  in Figure 3 is  $\mu$ -critical and non- $\alpha$ -critical, while the edge  $e$  of the graph  $G_2$  in the same figure is  $\alpha$ -critical and non- $\mu$ -critical.

Nevertheless, for König-Egerváry graphs and especially for bipartite graphs, there is a closed relationship between these two kinds of edges.

**Lemma 2.1** [11] *In a König-Egerváry graph,  $\alpha$ -critical edges are also  $\mu$ -critical, and they coincide in a bipartite graph.*




Figure 3: Non-König-Egervary graphs.

In a König-Egervary graph, maximum matchings have a very specific property, emphasized by the following statement:

**Lemma 2.2** [9] *Any maximum matching  $M$  of a König-Egervary graph  $G$  is contained in each  $(S, V(G) - S)$  and  $|M| = |V(G) - S|$ , where  $S \in \Omega(G)$ .*

Clearly, not any matching of a graph is contained in a maximum matching. For example, there is no maximum matching of the graph  $G$  in Figure 2 that includes the matching  $M = \{ab, cf\}$ . Let us observe that  $M$  is a maximum matching in  $G[N[\{a, f\}]]$ ,  $\{a, f\}$  is stable in  $G$ , but  $\{a, f\} \notin \Psi(G)$ . The following result shows that, under certain conditions, a matching of a bipartite graph can be extended to a maximum matching.

**Lemma 2.3** *If  $G$  is a bipartite graph,  $\widehat{S} \in \Psi(G)$ , and  $\widehat{M}$  is a maximum matching in  $G[N[\widehat{S}]]$ , then there exists a maximum matching  $M$  in  $G$  such that  $\widehat{M} \subseteq M$ .*

**Proof.** Let  $W = N(\widehat{S})$ ,  $H = G[N[\widehat{S}]]$ , and  $S'$  be a stable set in  $G$  such that  $S = \widehat{S} \cup S' \in \Omega(G)$  (such  $S'$  exists according to Theorem 1.1). Since  $H$  is bipartite and  $\widehat{M}$  is a maximum matching in  $H$ , it follows that

$$|\widehat{S}| + |\widehat{M}| = \alpha(H) + \mu(H) = |V(H)| = |\widehat{S}| + |W|.$$

Let  $M$  be a maximum matching in  $G$ . Then, by Lemma 2.2,  $M \subseteq (S, V(G) - S)$ , because  $S \in \Omega(G)$ , and

$$|M| = |V(G) - S| = |N(\bar{S})| + |N(S') - N(\widehat{S})| = |\widehat{M}| + |V(G) - S - W|.$$

Let  $M'$  be the subset of  $M$  containing edges having an endpoint in  $V(G) - S - W$ . Since no edge joins a vertex of  $\widehat{S}$  to some vertex in  $V(G) - S - W$ , it follows that  $M'$  is the restriction of  $M$  to  $G[V(G) - S - W]$ . Consequently,  $\widehat{M} \cup M'$  is a matching in  $G$  that contains  $\widehat{M}$ , and because  $|\widehat{M} \cup M'| = |\widehat{M}| + |V(G) - S - W| = |M|$ , we see that  $\widehat{M} \cup M'$  is a maximum matching in  $G$ . ■

Let us notice that Lemma 2.3 can not be generalized to non-bipartite graphs. For instance, the graph  $G$  presented in Figure 4 has  $\widehat{S} = \{a, d\} \in \Psi(G)$ ,  $\widehat{M} = \{ac, df\}$  is a maximum matching in  $G[N[\widehat{S}]]$ , but there is no maximum matching in  $G$  that includes  $\widehat{M}$ .




Figure 4:  $\widehat{M} = \{ac, df\}$  is a maximum matching in  $G[N[\{a, d\}]]$ .

**Lemma 2.4** *If  $G = (A, B, E)$  is a connected bipartite graph having a unique perfect matching, then  $A \cap \text{pend}(G) \neq \emptyset$  and  $B \cap \text{pend}(G) \neq \emptyset$ .*

**Proof.** Let  $M = \{a_i b_i : 1 \leq i \leq n, a_i \in A, b_i \in B\}$  be the unique perfect matching of  $G$ . Clearly,  $|A| = |B|$ . Suppose that  $B \cap \text{pend}(G) = \emptyset$ . Hence,  $|N(b_i)| \geq 2$  for any  $b_i \in B$ .

Under these conditions, we shall build some cycle  $C$  having half of edges contained in  $M$ , and this allows us to find a new perfect matching in  $G$ , which contradicts the uniqueness of  $M$ . We begin with the edge  $a_1 b_1$ . Since  $|N(b_1)| \geq 2$ , there is some  $a \in (A - \{a_1\}) \cap N(b_1)$ , say  $a_2$ . We continue with  $a_2 b_2 \in M$ . Further,  $N(b_2)$  contains some  $a \in (A - \{a_2\})$ . If  $a_1 \in N(b_2)$ , we are done, because  $G[\{a_1, a_2, b_1, b_2\}] = C_4$ . Otherwise, we may suppose that  $a = a_3$ , and we add to the growing cycle the edge  $a_3 b_3$ . Since  $G$  has a finite number of vertices, after a number of edges from  $M$ , we must find some edge  $a_j b_k$  with  $1 \leq j < k$ . So, the cycle  $C$  we found has

$$V(C) = \{a_i, b_i : j \leq i \leq k\}, \quad E(C) = \{a_i b_i : j \leq i \leq k\} \cup \{b_i a_{i+1} : j \leq i < k\} \cup \{a_j b_k\}.$$

Clearly, half of edges of  $C$  are contained in  $M$ .

Similarly, we can show that also  $A \cap \text{pend}(G) \neq \emptyset$ . ■

The following proposition presents a recursive structure of bipartite graphs owing unique perfect matchings, which generalizes the recursive structure of trees having perfect matching due to Fricke, Hedetniemi, Jacobs and Trevisan, [3].

**Proposition 2.5**  *$K_2$  is a bipartite graph, and it has a unique perfect matching. If  $G$  is a bipartite graph with a unique perfect matching, then  $G + K_2$  is also a bipartite graph having a unique perfect matching. Moreover, any bipartite graph containing a unique perfect matching can be obtained in this way.*

By  $G + K_2$  we mean the graph comprising the disjoint union of  $G$  and  $K_2$ , and additional edges joining at most one of endpoints of  $K_2$  to vertices belonging to only one color class of  $G$ .

**Proof.** Let  $G = (A, B, E)$  be a bipartite graph having a unique perfect matching, say  $M = \{a_i b_i : 1 \leq i \leq n, a_i \in A, b_i \in B\}$ . If  $K_2 = (\{x, y\}, \{xy\})$ , then  $H = G + K_2$  is also bipartite and  $M \cup \{xy\}$  is a unique perfect matching in  $H$ , since  $M$  was unique in  $G$  and at least one of  $x, y$  is pendant in  $H$ .

Conversely, let  $G$  be a bipartite graph with a unique perfect matching. By Lemma 2.4, it follows that  $G$  has at least one pendant vertex, say  $x$ . If  $y \in N(x)$ , then, clearly,  $G = (G - \{x, y\}) + K_2$ . ■

### 3 Main results

**Proposition 3.1** *If  $G$  is a bipartite graph of order  $2n$  having a perfect matching  $M$ , then  $M$  is unique if and only if for some  $S \in \Omega(G)$  there exists an accessibility chain.*

**Proof.** Since  $\mu(G) = n$ , in every set of size greater than  $n$  there exists a pair of adjacent vertices, and hence  $\alpha(G) = n$ .

Suppose that  $G$  is a bipartite graph of order  $2n$  with a unique perfect matching. We prove, by induction on  $n$ , that for some  $S \in \Omega(G)$  there exists an accessibility chain.

For  $n = 2$ , let  $S = \{x_1, x_2\} \in \Omega(G)$ ,  $N(S) = \{y_1, y_2\}$  and  $x_1y_1, x_2y_2 \in M$ , where  $M$  is its unique perfect matching. Then, at least one of  $x_1, x_2$  is pendant, say  $x_1$ . Hence,  $\{x_1\} \subset \{x_1, x_2\} = S$  is an accessibility chain.

Suppose that the assertion is true for  $k < n$ . Let  $G = (A, B, E)$  be of order  $2n$  and  $M = \{a_i b_i : 1 \leq i \leq n, a_i \in A, b_i \in B\}$  be its unique perfect matching. According to Proposition 2.5,  $G = H + K_2$ . Consequently, we may assume that:  $K_2 = (\{a_1, b_1\}, \{a_1 b_1\})$  and  $a_1 \in \text{pend}(G)$ . Clearly,  $H$  is a bipartite graph containing a unique perfect matching, namely  $M_H = M - \{a_1 b_1\}$ .

*Case 1.*  $a_1 \in S$ . Hence,  $S_{n-1} = S - \{a_1\} \in \Omega(H)$ , and by induction hypothesis, there is a chain

$$\{x_1\} \subset \{x_1, x_2\} \subset \dots \subset \{x_1, x_2, \dots, x_{n-2}\} \subset \{x_1, x_2, \dots, x_{n-1}\} = S_{n-1}$$

such that  $\{x_1, x_2, \dots, x_k\} \in \Psi(H)$  for any  $k \in \{1, \dots, n-1\}$ . Since  $N(a_1) = \{b_1\}$ , it follows that  $N_G(\{x_1, x_2, \dots, x_k\} \cup \{a_1\}) = N_H(\{x_1, x_2, \dots, x_k\}) \cup \{b_1\}$ , and therefore  $\{x_1, x_2, \dots, x_k\} \cup \{a_1\} \in \Psi(G)$  for any  $k \in \{1, \dots, n-1\}$ . Clearly,  $\{a_1\} \in \Psi(G)$ , and consequently, we have the chain:

$$\begin{aligned} \{a_1\} &\subset \{a_1, x_1\} \subset \{a_1, x_1, x_2\} \subset \dots \subset \{a_1, x_1, x_2, \dots, x_{n-2}\} \subset \\ &\subset \{a_1, x_1, x_2, \dots, x_{n-1}\} = \{a_1\} \cup S_{n-1} = S, \end{aligned}$$

where  $\{a_1, x_1, x_2, \dots, x_k\} \in \Psi(G)$ , for all  $k \in \{1, \dots, n-1\}$ .

*Case 2.*  $b_1 \in S$ . Hence,  $S_{n-1} = S - \{b_1\} \in \Omega(H)$  and also  $S_{n-1} \in \Psi(G)$ , because  $N_G[S_{n-1}] = A \cup B - \{a_1, b_1\}$ . By induction hypothesis, there is a chain

$$\{x_1\} \subset \{x_1, x_2\} \subset \dots \subset \{x_1, x_2, \dots, x_{n-2}\} \subset \{x_1, x_2, \dots, x_{n-1}\} = S_{n-1}$$

such that  $\{x_1, x_2, \dots, x_k\} \in \Psi(H)$  for any  $k \in \{1, \dots, n-1\}$ . Since none of  $a_1, b_1$  is contained in  $N_G(\{x_1, x_2, \dots, x_k\})$ , it follows that  $\{x_1, x_2, \dots, x_k\} \in \Psi(G)$ , for any  $k \in \{1, \dots, n-1\}$ . Consequently, we have the chain

$$\{x_1\} \subset \{x_1, x_2\} \subset \dots \subset \{x_1, x_2, \dots, x_{n-1}\} = S_{n-1} \subset S_{n-1} \cup \{b_1\} = S,$$

where  $\{x_1, x_2, \dots, x_k\} \in \Psi(G)$ , for all  $k \in \{1, \dots, n-1\}$ .

Conversely, let  $M = \{x_i y_i : 1 \leq i \leq n\}$  be a perfect matching in  $G$ , and suppose that for  $S \in \Omega(G)$  there exists a chain of local maximum stable sets

$$\{x_1\} \subset \{x_1, x_2\} \subset \dots \subset \{x_1, x_2, \dots, x_{n-1}\} \subset \{x_1, x_2, \dots, x_{\alpha-1}, x_{\alpha}\} = S.$$

We show, by induction on  $k = |\{x_1, x_2, \dots, x_k\}|$  that  $H_k = G[N[\{x_1, x_2, \dots, x_k\}]]$  owns a unique perfect matching.

For  $k = 1$ , the assertion is true, because  $\{x_1\} \in \Psi(G)$  ensures that  $x_1$  is pendant, and therefore,  $H_1 = G[N[\{x_1\}]]$  has a unique perfect matching, consisting of the unique edge issuing from  $x_1$ , namely  $x_1y_1$ .

Assume that  $H_k$  has a unique perfect matching, say  $M_k$ . We may assert that  $M_k \subseteq M$ , because  $M_k$  is unique and included in  $H_k$  and also  $M$  matches  $x_1, x_2, \dots, x_k$  onto vertices belonging to  $N(\{x_1, x_2, \dots, x_k\})$ . Hence,  $M_{k+1} = M_k \cup \{x_{k+1}y_{k+1}\}$  is a maximum matching in  $H_{k+1}$ . If  $M_{k+1}$  is not unique in  $H_{k+1}$ , then there exists some  $z \in N(a_{k+1}) - N[\{a_1, a_2, \dots, a_k\}]$  such that  $z \neq y_{k+1}$ . Therefore, we infer that the set  $\{x_1, x_2, \dots, x_k\} \cup \{z, y_{k+1}\}$  is stable in  $H_{k+1}$  and larger than  $\{x_1, x_2, \dots, x_{k+1}\}$ , which contradicts the fact that  $\{x_1, x_2, \dots, x_{k+1}\} \in \Psi(G)$ . Consequently,  $M_{k+1}$  is unique and also perfect in  $H_{k+1}$ . ■

If one of the maximum matchings of a bipartite graph is uniquely restricted, this is not necessarily true for all its maximum matchings. For instance, let us consider the bipartite graph  $G$  presented in Figure 5. The set of edges  $M_1 = \{ab, ce\}$  is one of uniquely restricted maximum matchings of  $G$ , while  $M_2 = \{bd, cf\}$  is one of its maximum matchings, but it is not uniquely restricted.




Figure 5: Not all maximum matchings of a graph have to be uniquely restricted.

**Theorem 3.2** *If  $G$  is a bipartite graph, then the following assertions are equivalent:*

- (i) *there exists some  $S \in \Omega(G)$  having an accessibility chain;*
- (ii) *there exists a uniquely restricted maximum matching in  $G$ ;*
- (iii) *each  $S \in \Omega(G)$  has an accessibility chain.*

**Proof.** (i)  $\Rightarrow$  (ii) Let us consider an accessibility chain of  $S \in \Omega(G)$

$$\emptyset \subset \{x_1\} \subset \{x_1, x_2\} \subset \dots \subset \{x_1, x_2, \dots, x_{\alpha-1}\} \subset \{x_1, x_2, \dots, x_\alpha\} = S,$$

for which we define  $S_i = \{x_1, x_2, \dots, x_i\}$  and  $S_0 = \emptyset$ .

Since  $S_{i-1} \in \Psi(G)$ ,  $S_i = S_{i-1} \cup \{x_i\} \in \Psi(G)$  and  $G$  is bipartite, it follows that  $|N(x_i) - N[S_{i-1}]| \leq 1$ , because otherwise, if  $\{a, b\} \subset N(x_i) - N[S_{i-1}]$ , then the set  $\{a, b\} \cup S_{i-1}$  is stable in  $N[S_{i-1} \cup \{x_i\}]$ , and larger than  $S_i = S_{i-1} \cup \{x_i\}$ , in contradiction with the fact that  $S_i \in \Psi(G)$ .

Let  $\{y_{i_j} : 1 \leq j \leq \mu\}$  be such that  $\{y_{i_j}\} = N(x_{i_j}) - N[S_{i_j-1}]$ , for all  $i \in \{1, \dots, \alpha\}$  with  $|N(x_i) - N[S_{i-1}]| = 1$ . Hence,  $M = \{x_{i_j}y_{i_j} : 1 \leq j \leq \mu\}$  is a matching in  $G$ .

- *Claim 1.*  $\mu = \mu(G)$ , i.e.,  $M$  is a maximum matching in  $G$ .

Since  $|N(x_i) - N[S_{i-1}]| \leq 1$  holds for all  $i \in \{1, \dots, \alpha\}$ , where  $S_0 = N[S_0] = \emptyset$ , and  $\{y_{i_j}\} = N(x_{i_j}) - N[S_{i_j-1}]$ , for all  $i \in \{1, \dots, \alpha\}$  satisfying  $|N(x_i) - N[S_{i-1}]| = 1$ , it follows that  $N(S) = \{y_{i_j} : 1 \leq j \leq \mu\}$ , and this ensures that  $M$  is a maximal matching in  $G$ , i.e., it is impossible to add an edge to  $M$  and to get a new matching.

In addition, we have

$$|V(G)| = |N[S]| = |S| + |N(S)| = |S| + |\{y_{i_j} : 1 \leq j \leq \mu\}| = \alpha(G) + |M|,$$

and because  $|V(G)| = \alpha(G) + \mu(G)$ , we infer that  $|M| = \mu(G)$ . In other words,  $M$  is a maximum matching in  $G$ .

- *Claim 2.*  $M$  is a uniquely restricted maximum matching in  $G$ .

We use induction on  $k = |S_k|$  to show that the restriction of  $M$  to  $H_k = G(N[S_k])$ , which we denote by  $M_k$ , is a uniquely restricted maximum matching in  $H_k$ .

For  $k = 1$ ,  $S_1 = \{x_1\} \in \Psi(G)$  and this implies that  $N(x_1) = \{y_{i_1}\}$ . Clearly,  $M_1 = \{x_1 y_{i_1}\}$  is a uniquely restricted maximum matching in  $H_1$ .

Suppose that the assertion is true for all  $j \leq k - 1$ . Let us observe that

$$N[S_k] = N[S_{k-1}] \cup (N(x_k) - N[S_{k-1}]) \cup \{x_k\},$$

because  $S_k = S_{k-1} \cup \{x_k\}$ .

Further we will distinguish between two different situations depending on the number of new vertices, which the set  $N(x_k)$  brings to the set  $N[S_{k-1}]$ .

*Case 1.*  $N(x_k) - N[S_{k-1}] = \emptyset$ . Hence, we obtain:

$$|V(H_k)| = |S_{k-1} \cup \{x_k\}| + |M_{k-1}| = |S_k| + |M_{k-1}| = \alpha(H_k) + |M_{k-1}|.$$

Since  $|V(H_k)| = \alpha(H_k) + \mu(H_k)$ , the equality  $|V(H_k)| = \alpha(H_k) + |M_{k-1}|$  ensures that  $M_{k-1}$  is a maximum matching of  $H_k$ . Therefore,  $M_{k-1}$  is a uniquely restricted maximum matching in  $H_k$ .

*Case 2.*  $N(x_k) - N[S_{k-1}] = \{y_{i_k}\}$ . Then we have:

$$|V(H_k)| = |S_{k-1} \cup \{x_k\}| + |M_{k-1} \cup \{x_k y_{i_k}\}| = |S_k| + |M_k| = \alpha(H_k) + |M_k|,$$

and this assures that  $M_k = M_{k-1} \cup \{x_k y_{i_k}\}$  is a maximum matching in  $H_k$ . The edge  $e = x_k y_{i_k}$  is  $\alpha$ -critical in  $H_k$ , because  $\{y_{i_k}\} = N(x_k) - N[S_{k-1}]$ . According to Lemma 2.1,  $e$  is also  $\mu$ -critical in  $H_k$ . Therefore, any maximum matching of  $H_k$  contains  $e$ , and since  $M_k = M_{k-1} \cup \{e\}$  and  $M_{k-1}$  is a uniquely restricted maximum matching in  $H_{k-1} = H_k - \{x_k, y_{i_k}\}$ , it follows that  $M_k$  is a uniquely restricted maximum matching in  $H_k$ .

(ii)  $\Rightarrow$  (iii) Let  $M$  be a uniquely restricted maximum matching in  $G$ . According to Lemma 2.2,  $M \subseteq (S, V(G) - S)$  and  $|M| = |V(G) - S| = \mu(G)$ . Therefore,  $M$  is a unique perfect matching in  $H = G[N[S_\mu]]$ , where

$$S_\mu = \{x : x \in S, x \text{ is an endpoint of an edge in } M\}.$$



It is clear that  $S_\mu$  is a maximum stable set in  $H$ , because  $N(S_\mu) = V(G) - S$  and  $S_\mu$  is stable. In other words,  $S_\mu \in \Psi(G)$ . Since  $H$  is bipartite and  $M$  is its unique perfect matching, Proposition 3.1 implies that there exists a chain

$$\{x_1\} \subset \{x_1, x_2\} \subset \dots \subset \{x_1, x_2, \dots, x_{\mu-1}\} \subset \{x_1, x_2, \dots, x_{\mu-1}, x_\mu\} = S_\mu,$$

such that all  $S_k = \{x_1, x_2, \dots, x_k\}$ ,  $1 \leq k \leq \mu$  are local maximum stable sets in  $H$ . The equality  $N_H[S_k] = N_G[S_k]$  explains why  $S_k \in \Psi(G)$  for all  $k \in \{1, \dots, \mu(G)\}$ . Let now  $x \in S - S_\mu$ . Then  $N(x) \subseteq V(G) - S$ , and therefore,  $N(S_\mu \cup \{x\}) = V(G) - S$ . Since  $S_\mu$  is a maximum stable set in  $H$  and  $S_\mu \cup \{x\}$  is stable in  $H \cup \{x\} = G[N[S_\mu \cup \{x\}]]$ , we get that  $S_\mu \cup \{x\}$  is a maximum stable set in  $H \cup \{x\}$ , i.e.,  $S_{\mu+1} = S_\mu \cup \{x\} \in \Psi(G)$ . If there still exists some  $y \in S - S_{\mu+1}$ , in the same manner as above we infer that  $S_{\mu+2} = S_{\mu+1} \cup \{y\} \in \Psi(G)$ .

In such a way we build the following accessibility chain

$$\{x_1\} \subset \{x_1, x_2\} \subset \dots \subset \{x_1, x_2, \dots, x_\mu\} \subset S_{\mu+1} \subset S_{\mu+1} \subset \dots \subset S_\alpha = S.$$

Clearly, (iii)  $\Rightarrow$  (i), and this completes the proof. ■

As an example of the process of building a uniquely restricted maximum matching with the help of an accessibility chain, let us consider the bipartite graph  $G$  presented in Figure 6. The accessibility chain

$$\{h\} \subset \{h, d\} \subset \{h, d, f\} \subset \{h, d, f, c\} \subset \{h, d, f, c, a\} \in \Psi(G)$$

gives rise to the uniquely restricted maximum matching  $M = \{hg, de, cb\}$ . Notice that  $\Psi(G)$  is not a greedoid, because  $\{d, f\} \in \Psi(G)$ , while  $\{d\}, \{f\} \notin \Psi(G)$ .




Figure 6: The chain of uniquely restricted matchings is  $\{\{hg\}, \{hg, de\}, \{hg, de, cb\}\}$ .

The following theorem will show us another reason, why the family  $\Psi(G)$  of the graph  $G$  from Figure 6 is not a greedoid, namely  $\{bc, de, fg\}$  is a maximum matching, but not uniquely restricted.

**Theorem 3.3** *If  $G$  is a bipartite graph, then  $\Psi(G)$  is a greedoid if and only if all its maximum matchings are uniquely restricted.*

**Proof.** Assume that  $\Psi(G)$  is a greedoid. Let  $M$  be a maximum matching in  $G$ . According to Lemma 2.2, we have that  $M \subseteq (S, V(G) - S)$  and  $|M| = |V(G) - S|$  for any  $S \in \Omega(G)$ . Let  $S_\mu$  contain the vertices of some  $S \in \Omega(G)$  matched by  $M$  with the vertices of  $V(G) - S$ . Since  $M$  is a perfect matching in  $G[N[S_\mu]]$  and  $|S_\mu| = |M|$ , it follows that  $S_\mu$  is a maximum stable set in  $G[N[S_\mu]]$ , i.e.,  $S_\mu \in \Psi(G)$ . Hence, there exists an accessibility chain of the following structure:

$$\{x_1\} \subset \{x_1, x_2\} \subset \dots \subset \{x_1, x_2, \dots, x_\mu\} = S_\mu \subset S_\mu \cup \{x_{\mu+1}\} \subset \dots \subset S.$$

While the existence of the first part of this chain, i.e.,  $\{x_1\}, \{x_1, x_2\}, \dots, \{x_1, x_2, \dots, x_\mu\}$ , is based on the accessibility property of the family  $\Psi(G)$ , the existence of the second part of the same chain, namely  $S_\mu, S_\mu \cup \{x_{\mu+1}\}, \dots, S$ , stems from the exchange property of  $\Psi(G)$ . Now, according to Proposition 3.1, we may conclude that the perfect matching  $M$  is unique in  $G[N[S_\mu]]$ . Hence,  $M$  is a uniquely restricted maximum matching in  $G$ .

Conversely, suppose that all maximum matchings of  $G$  are uniquely restricted. Let  $\hat{S} \in \Psi(G)$ ,  $H = G[N[\hat{S}]]$ , and  $\hat{M}$  be a maximum matching in  $H$ . The graph  $H$  is bipartite as a subgraph of a bipartite graph. By Lemma 2.3, there exists a maximum matching in  $G$ , say  $M$ , such that  $\hat{M} \subseteq M$ . Since  $M$  is uniquely restricted in  $G$ , it follows that  $\hat{M}$  is uniquely restricted in  $H$ . According to Theorem 3.2, there exists an accessibility chain of  $\hat{S}$  in  $H$

$$S_1 \subset S_2 \subset \dots \subset S_{q-1} \subset S_q = \hat{S}.$$

Since  $N_H[S_k] = N_G[S_k]$ , we infer that  $S_k \in \Psi(G)$ , for any  $k \in \{1, \dots, q\}$ .

To complete the proof, we have to show that, in addition to the accessibility property,  $\Psi(G)$  satisfies also the exchange property.

Let  $X, Y \in \Psi(G)$  and  $|Y| = |X| + 1 = m + 1$ . Hence, there is an accessibility chain

$$\{y_1\} \subset \{y_1, y_2\} \subset \dots \subset \{y_1, \dots, y_m\} \subset \{y_1, \dots, y_m, y_{m+1}\} = Y.$$

Since  $Y$  is stable,  $X \in \Psi(G)$ , and  $|X| < |Y|$ , it follows that there exists some  $y \in Y - X$ , such that  $y \notin N[X]$ . Let  $M_X$  be a maximum matching in  $H = G[N[X]]$ . Since  $H$  is bipartite,  $X$  is a maximum stable set in  $H$ , and  $M_X$  is a maximum matching in  $H$ , it follows that

$$|X| + |M_X| = |N[X]| = |X| + |N(X)|, \text{ i.e., } |M_X| = |N(X)|.$$

Let  $y_{k+1} \in Y$  be the first vertex in  $Y$  satisfying the conditions:  $y_1, \dots, y_k \in N[X]$  and  $y_{k+1} \notin N[X]$ . Since  $\{y_1, \dots, y_k\}$  is stable in  $N[X]$ , there is  $\{x_1, \dots, x_k\} \subseteq X$  such that for any  $i \in \{1, \dots, k\}$  either  $x_i = y_i$  or  $x_i y_i \in M_X$ .

Now we show that  $X \cup \{y_{k+1}\} \in \Psi(G)$ .

*Case 1.*  $N[X \cup \{y_{k+1}\}] = N[X] \cup \{y_{k+1}\}$ . Clearly,  $X \cup \{y_{k+1}\}$  is stable in  $G(N[X \cup \{y_{k+1}\}])$  and  $|X \cup \{y_{k+1}\}| = |X| + 1$  ensures that  $X \cup \{y_{k+1}\} \in \Psi(G)$ , because  $X \in \Psi(G)$  too.

*Case 2.*  $N[X \cup \{y_{k+1}\}] \neq N[X] \cup \{y_{k+1}\}$ . Suppose there are  $a, b \in N(y_{k+1}) - N[X]$ . Hence, it follows that  $\{a, b, x_1, \dots, x_k\}$  is a stable set included in  $N[\{y_1, \dots, y_{k+1}\}]$  and larger than  $\{y_1, \dots, y_{k+1}\}$ , in contradiction with the fact that  $\{y_1, \dots, y_{k+1}\} \in \Psi(G)$ . Therefore, there exists a unique  $a \in N(y_{k+1}) - N[X]$ . Consequently,

$$N[X \cup \{y_{k+1}\}] = N[X] \cup N[y_{k+1}] = N[X] \cup \{a, y_{k+1}\},$$

and since  $ay_{k+1} \in E(G)$ , we obtain that  $X \cup \{y_{k+1}\}$  is a maximum stable set in  $G[N[X \cup \{y_{k+1}\}]]$ , i.e.,  $X \cup \{y_{k+1}\} \in \Psi(G)$ . ■

As an immediate consequence of Theorem 3.3, we obtain the following:

**Corollary 3.4** *For any bipartite graph  $G$  having a perfect matching,  $\Psi(G)$  is a greedoid if and only if  $G$  has a unique perfect matching.*

Corollary 3.4 and, consequently, Theorem 3.3 are not valid for non-bipartite graphs. For example, the graph  $C_5 + e$  in Figure 7 is a non-bipartite graph having only uniquely restricted maximum matchings, (in fact, it has a unique perfect matching), but  $\Psi(C_5 + e)$  is not a greedoid, because  $\{u, v\} \in \Psi(C_5 + e)$ , while  $\{u\}, \{v\} \notin \Psi(C_5 + e)$ .




Figure 7: Non-bipartite graphs with unique perfect matchings.

However, there are non-bipartite graphs with a unique perfect matching, whose  $\Psi(G)$  is a greedoid. For instance, while the graph  $C_5 + 3e$  in Figure 7 is a non-bipartite graph with a unique perfect matching, the family  $\Psi(C_5 + 3e)$  is a greedoid.

Let us also notice that there exist both bipartite and non-bipartite graphs without a perfect matching whose family of local maximum stable sets is a greedoid. For instance, neither  $G_1$  nor  $G_2$  in Figure 8 has a perfect matching,  $G_1$  is bipartite, and  $\Psi(G_1), \Psi(G_2)$  are greedoids.



Figure 8:  $\Psi(G_1)$  and  $\Psi(G_2)$  form greedoids, but only  $G_1$  is a bipartite graph.

Since any forest, by definition, has no cycles, the following Lemma 3.5 ensures that all matchings of a forest are uniquely restricted.

**Lemma 3.5** [8] *If a bipartite graph has two perfect matchings  $M_1$  and  $M_2$ , then any of its vertices, from which are issuing edges contained in  $M_1$  and  $M_2$ , respectively, belongs to some cycle that is alternating with respect to at least one of  $M_1, M_2$ .*

It is also interesting to note that Golumbic, Hirst and Lewenstein have proved the following generalization of Lemma 3.5.

**Theorem 3.6** [4] *A matching  $M$  in a graph  $G$  is uniquely restricted if and only if there is no even-length cycle with edges alternating between matched and non-matched edges.*

Now restricting Theorem 3.3 to forests we immediately obtain that the family of local maximum stable sets of a forest forms a greedoid on its vertex set, which gives a new proof of the main finding from [10], namely Theorem 1.2.

## 4 Conclusions

We have shown that to have all maximum matchings uniquely restricted is necessary and sufficient for a bipartite graph  $G$  to enjoy the property that  $\Psi(G)$  is a greedoid. We have also described all the bipartite graphs having a unique perfect matching, or in other words, all bipartite graphs having a perfect matching and whose  $\Psi(G)$  is a greedoid. It seems to be interesting to describe a recursive structure of all bipartite graphs whose  $\Psi(G)$  is a greedoid.

A linear time algorithm to decide whether a matching in a bipartite graph is uniquely restricted is presented in [4]. It is also shown there that the problem of finding a maximum uniquely restricted matching is **NP**-complete for bipartite graphs. These results motivate us to propose another open problem, namely: how to recognize bipartite graphs whose  $\Psi(G)$  is a greedoid?

## References

- [1] A. Björner and G. M. Ziegler, *Introduction to greedoids*, in N. White (ed.), *Matroid Applications*, 284-357, Cambridge University Press, 1992.
- [2] E. Egervary, *On combinatorial properties of matrices*, Mat. Lapok **38** (1931) 16-28.
- [3] G. H. Fricke, S. T. Hedetniemi, D. P. Jacobs and V. Trevisan, *Reducing the adjacency matrix of a tree*, The Electronic Journal of Linear Algebra, vol. **1** (1996) 34-43.
- [4] M. C. Golumbic, T. Hirst and M. Lewenstein, *Uniquely restricted matchings*, Algorithmica (2000).
- [5] D. Hershkowitz and H. Schneider, *Ranks of zero patterns and sign patterns*, Linear and Multilinear Algebra **34** (1993) 3-19.
- [6] B. Korte, L. Lovász and R. Schrader, *Greedoids*, Springer-Verlag, Berlin, 1991.
- [7] D. König, *Graphen und Matrizen*, Mat. Lapok **38** (1931) 116-119.
- [8] V. E. Levit and E. Mandrescu, *The structure of  $\alpha$ -stable graphs*, Discrete Mathematics (2000) (accepted).
- [9] V. E. Levit and E. Mandrescu, *On  $\alpha^+$ -stable König-Egervary graphs*, Los Alamos Archive, pre-print arXiv:math.CO/9912022, 1999, 13 pp.
- [10] V. E. Levit and E. Mandrescu, *A new greedoid: the family of local maximum stable sets of a forest*, Discrete Applied Mathematics (2000). (accepted)
- [11] V. E. Levit and E. Mandrescu, *On  $\alpha$ -critical edges in König-Egervary graphs*, Los Alamos Archive, pre-print arXiv:math.CO/0002070, 2000, 15 pp.
- [12] G. L. Nemhauser and L. E. Trotter, Jr., *Vertex packings: structural properties and algorithms*, Mathematical Programming **8** (1975) 232-248.